arXiv:1112.5811v2 [math.AT] 26 Jan 2012

COHOMOLOGY MOD 3 OF THE CLASSIFYING SPACE OF THE EXCEPTIONAL LIE GROUP E_6, I : STRUCTURE OF COTOR

MAMORU MIMURA, YURIKO SAMBE, AND MICHISHIGE TEZUKA

Abstract. We study the structure of the E_2-term of the Rothenberg-Steenrod spectral sequence converging to the classifying space of the compact, connected, simply connected, exceptional Lie group of rank 6.

2010 Mathematics Subject Classification. 55R35, 55R40, 55T20

Key words and phrases, exceptional Lie group, spectral sequence, Cotor

1. Introduction

Let E_6 be the compact, connected, simply connected, exceptional Lie group of rank 6 and BE_6 its classifying space. Then we have the Rothenberg-Steenrod spectral sequence $\{E_r, d_r\}$ ([RS]) such that

$$E_2 = \mathrm{Cotor}_{H^*(E_6;\mathbb{Z}_3)}(\mathbb{Z}_3, \mathbb{Z}_3) \quad \text{and} \quad E_\infty = \mathrm{gr}\, H^*(BE_6; \mathbb{Z}_3).$$

The E_2-term was calculated by Mimura and Sambe [MS], where some details of the calculations are somewhat omitted. This is one of the reasons why we give here a detailed account of them. After the calculations of the E_2-term, the collapsing of the spectral sequence was proved by Kono and Mimura [KM]; thus one can obtain the module structure of $H^*(BE_6; \mathbb{Z}_3)$. (See also Toda [T], in which the collapsing is shown by a different method.) Further in order to determine the ring structure of $H^*(BE_6; \mathbb{Z}_3)$, we will need more information (see [MST1]), that is, we need to add supplement to the results of [MS]. This is the main reason for the present work.

The paper is organized as follows. In Section 2 we state our results. In Section 3 we prove Theorem 2.1 and in Section 4 we prove Theorems 2.3.A and 2.3.B. In Section 5, we compare our calculation to the May spectral sequence.

Thus the present work is essentially a supplement, and hence a more detailed version, of [KM] and [MS]. The results of the present work are announced in [MST2].

In the series of our work, we will determine $H^*(BE_6; \mathbb{Z}_3)$. Continuation follows.

Throughout the paper, the coefficient field is always $\mathbb{Z}_3$, which is hereafter omitted.

Partially supported by the Grant-in-Aid for Scientific Research (C) 21540104, Japan Society for the Promotion of Science.

2. Results

Let us start with

Notation For a commutative ring R, we denote by $R\{x_1, \dots, x_n\}$ a free R-module generated by elements $x_1, \dots, x_n$. For a graded algebra S, we denote by S^+ the ideal generated by the elements with positive degrees. We also denote by $T(x_1, \dots, x_n)$ a tensor algebra generated by elements $x_1, \dots, x_n$, and by $\Delta(x)$ an algebra additively isomorphic to $\mathbb{Z}_3 \oplus \mathbb{Z}_3\{x\}$.

The following is essentially the restatement of Theorem 5.20 of [MS].

Theorem 2.1. $\mathrm{Cotor}_{H^*(E_6)}(\mathbb{Z}_3, \mathbb{Z}_3)$ *is additively isomorphic to the following module:*

$$(C \oplus D) \otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}],$$

where
$$\begin{aligned} C &= \mathbb{Z}_3[a_4, a_8, a_{10}]\{1, y_{20}, y_{20}^2, y_{22}, y_{22}^2, y_{20}y_{22}, y_{58}, y_{60}, y_{76}\} \\ &\qquad \oplus \mathbb{Z}_3[a_8, a_{10}]\{y_{26}, y_{26}^2, y_{20}y_{26}, y_{22}y_{26}, y_{64}\} \end{aligned}$$
and
$$\begin{aligned} D &= \mathbb{Z}_3[x_{26}]^+\{1, a_4, a_8, a_{10}, y_{20}, y_{22}, a_{10}y_{20}, y_{26}\} \\ &\qquad \oplus \mathbb{Z}_3[x_{26}]\{a_9, y_{21}, y_{25}, y_{27}, y_{21}a_8, y_{21}a_{10}, y_{25}a_{10}, y_{21}y_{26}\}. \end{aligned}$$

By an easy calculation one obtains

Corollary 2.2. *The Poincaré polynomials PS of them are given respectively by*

$$PS(C \otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}]) = \frac{g(t)}{(1-t^4)(1-t^8)(1-t^{10})} \cdot \frac{1}{(1-t^{36})(1-t^{48})(1-t^{54})},$$

$$PS(D \otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}]) = \frac{h(t)}{1-t^{26}} \cdot \frac{1}{(1-t^{36})(1-t^{48})(1-t^{54})},$$

where
$$\begin{aligned} g(t) &= 1 + t^{20} + t^{22} + t^{26} - t^{30} + t^{40} + t^{42} + t^{44} + t^{46} + t^{48} - t^{50} - t^{56} + t^{58} \\ &\qquad + t^{60} + t^{64} - t^{68} + t^{76} \end{aligned}$$
and
$$\begin{aligned} h(t) &= t^9 + t^{21} + t^{25} + t^{26} + t^{27} + t^{29} + t^{30} + t^{31} + t^{34} + t^{35} + t^{36} + t^{46} + t^{47} \\ &\qquad + t^{48} + t^{52} + t^{56}. \end{aligned}$$

The following two theorems will play an important role when determining the ring structure of $H^*(BE_6)$ (see [MST1]).

Theorem 2.3.A *The extension*

$$0 \to D \otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}] \to \mathrm{Cotor}_{H^*(E_6)}(\mathbb{Z}_3, \mathbb{Z}_3) \to C \otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}] \to 0$$

is split as algebras.

Theorem 2.3.B *We have an algebra isomorphism*

$$\begin{aligned} \mathrm{Cotor}_{H^*(E_6)}(\mathbb{Z}_3, \mathbb{Z}_3) \cong \mathbb{Z}_3[&a_9, x_{26}, y_{21}, y_{25}, y_{27}, a_4, a_8, a_{10}, x_{36}, \\ &x_{48}, x_{54}, y_{20}, y_{22}, y_{26}, y_{58}, y_{60}, y_{64}, y_{76}]/I, \end{aligned}$$

where I is the ideal generated by the following i)

$$
\begin{aligned}
a_4y_{26} &= -a_8y_{22}+a_{10}y_{20},\\
a_4y_{64} &= -a_8y_{60}-a_8^3y_{22}^2-a_{10}y_{58},\\
y_{22}^3 &= -a_4^3x_{54}-a_4^2a_8^2a_{10}^2y_{22}+a_4a_8a_{10}y_{22}^2+a_4a_{10}^2y_{20}y_{22}+a_{10}^3x_{36},\\
y_{20}^2y_{22} &= -a_4y_{58}+a_8^2a_{10}x_{36},\\
y_{20}y_{22}^2 &= -a_4y_{60}+a_8a_{10}^2x_{36},\\
y_{58}y_{22} &= -a_4y_{76}-a_4a_8^2y_{60}+a_8^3a_{10}^2x_{36}+a_8a_{10}x_{36}y_{26},\\
y_{60}y_{22} &= a_4^2x_{54}y_{20}+a_4a_8^2a_{10}^2y_{20}y_{22}+a_4a_8a_{10}y_{60}+a_4a_{10}^2y_{58}\\
&\qquad +a_8^2a_{10}^3x_{36}-a_{10}^2x_{36}y_{26},\\
y_{76}y_{22} &= -a_4x_{54}y_{20}^2-a_4^2a_8^2x_{54}y_{20}+a_4^2a_{10}^2x_{48}y_{22}+a_4a_8^4a_{10}^2y_{20}y_{22}\\
&\qquad +a_4a_8^2a_{10}^2y_{58}+a_4a_8a_{10}y_{76}-a_8^4a_{10}^3x_{36}+a_{10}x_{36}y_{26}^2,\\
y_{22}y_{26}^2 &= -a_4a_8^2x_{54}-a_8^4a_{10}^2y_{22}+a_8^2a_{10}y_{22}y_{26}+a_{10}y_{64},\\
y_{20}y_{22}y_{26} &= a_8y_{60}-a_{10}y_{58},\\
y_{22}^2y_{26} &= a_4^2a_8x_{54}+a_4a_8^3a_{10}^2y_{22}-a_8^2a_{10}y_{22}^2-a_8a_{10}^2y_{20}y_{22}-a_{10}y_{60},\\
y_{64}y_{22} &= a_4^2a_8^3x_{54}-a_4a_8x_{54}y_{20}+a_4a_8^5a_{10}^2y_{22}-a_8^4a_{10}y_{22}^2+a_8^3a_{10}^2y_{20}y_{22}\\
&\qquad -a_8a_{10}^2y_{58}+a_{10}y_{76},\\
y_{20}^3 &= a_4^3x_{48}-a_4^2a_8^4y_{20}-a_4a_8^2y_{20}^2+a_8^3x_{36},\\
y_{20}^2y_{22} &= -a_4y_{58}+a_8^2a_{10}x_{36},\\
y_{58}y_{20} &= -a_4^2x_{48}y_{22}-a_4a_8^2y_{58}+a_4a_8^4y_{20}y_{22}+a_8^2x_{36}y_{26}+a_8^4a_{10}x_{36},\\
y_{60}y_{20} &= -a_4y_{76}-a_4a_8^2y_{60}+a_8^3a_{10}^2x_{36}-a_8a_{10}x_{36}y_{26},\\
y_{76}y_{20} &= a_4x_{48}y_{22}^2+a_4a_8^4y_{60}+a_8x_{36}y_{26}^2-a_8^5a_{10}^2x_{36},\\
y_{20}y_{26}^2 &= a_4a_{10}^2x_{48}+a_8y_{64}-a_8^3y_{22}y_{26}-a_8^4a_{10}^2y_{20}-a_8^2a_{10}y_{20}y_{26},\\
y_{20}^2y_{26} &= a_4^2a_{10}x_{48}-a_4a_8^4a_{10}y_{20}+a_8y_{58}-a_8^2a_{10}y_{20}^2,\\
y_{64}y_{20} &= a_4a_{10}x_{48}y_{22}+a_8y_{76}-a_8^3y_{60}-a_8^4a_{10}y_{20}y_{22}+a_8^2a_{10}y_{58},\\
y_{58}^2 &= -a_4^2x_{48}y_{60}-a_4^2a_8^2x_{48}y_{22}^2-a_4^2a_8^6y_{60}+a_4a_8^7a_{10}^2x_{36}+a_4a_8a_{10}^2x_{36}x_{48}+a_8^2x_{36}y_{64},\\
y_{58}y_{60} &= -a_4^4x_{48}x_{54}+a_4^3a_8^4x_{54}y_{20}+a_4^3a_8^2a_{10}^2x_{48}y_{22}+a_4^2a_8^2x_{54}y_{20}^2-a_4^2a_8^6a_{10}^2y_{20}y_{22}\\
&\qquad -a_4^2a_8^4a_{10}^2y_{58}-a_4^2a_8^3a_{10}y_{76}+a_4^2a_8a_{10}x_{48}y_{22}^2+a_4^2a_{10}^2x_{48}y_{20}y_{22}-a_4a_8^3x_{36}x_{54}\\
&\qquad +a_4a_8^6a_{10}^3x_{36}+a_4a_{10}^3x_{36}x_{48}-a_8^5a_{10}^2x_{36}y_{22}-a_8^2a_{10}^2x_{36}y_{20}y_{26}-a_8a_{10}x_{36}y_{64},\\
y_{58}y_{76} &= a_4^3x_{48}x_{54}y_{20}+a_4^3a_8^4a_{10}^2x_{48}y_{22}-a_4^2a_8^4x_{54}y_{20}^2-a_4^2a_8^8a_{10}^2y_{20}y_{22}+a_4^2a_8^7a_{10}y_{60}\\
&\qquad -a_4^2a_8^6a_{10}^2y_{58}+a_4^2a_8^5a_{10}y_{76}+a_4^2a_8^2a_{10}^2x_{48}y_{20}y_{22}+a_4^2a_8a_{10}x_{48}y_{60}\\
&\qquad +a_4^2a_{10}^2x_{48}y_{58}+a_4a_8^2a_{10}^3x_{36}x_{48}+a_8^3x_{36}x_{54}y_{20}-a_8^6a_{10}^3x_{36}y_{20}\\
&\qquad +a_8^4a_{10}^2x_{36}y_{20}y_{26}-a_8a_{10}^2x_{36}x_{48}y_{22}-a_{10}^3x_{36}x_{48}y_{20},\\
y_{58}y_{26} &= -a_4a_{10}x_{48}y_{22}+a_8y_{76}+a_8^3y_{60}+a_8^4a_{10}y_{20}y_{22}-a_8^2a_{10}y_{58},\\
y_{58}y_{64} &= a_4^3a_8x_{48}x_{54}-a_4^2a_8^5x_{54}y_{20}+a_4a_8^3x_{54}y_{20}^2-a_4a_8^6a_{10}y_{60}-a_4a_8^4a_{10}y_{76}\\
&\qquad -a_4a_8a_{10}^2x_{48}y_{20}y_{22}+a_4a_{10}x_{48}y_{60}+a_8^4x_{36}x_{54}+a_8^7a_{10}^3x_{36}-a_8^5a_{10}^2x_{36}y_{26}\\
&\qquad -a_8^3a_{10}x_{36}y_{26}^2+a_8a_{10}^3x_{36}x_{48},
\end{aligned}
$$

$$
\begin{aligned}
y_{60}^2 &= a_4^2x_{54}y_{58}+a_4^3a_8a_{10}^3x_{48}y_{22}-a_4^2a_8^5a_{10}^3y_{20}y_{22}-a_4^2a_8^3a_{10}^3y_{58}-a_4^2a_8^2a_{10}^2y_{76}\\
&\qquad -a_4^2a_8a_{10}x_{54}y_{20}^2+a_4^2a_{10}^2x_{48}y_{22}^2+a_4a_8^5a_{10}^4x_{36}-a_4a_8^2a_{10}x_{36}x_{54}-a_8^4a_{10}^3x_{36}y_{22}\\
&\qquad +a_8^3a_{10}^4x_{36}y_{20}+a_8^2a_{10}^2x_{36}y_{22}y_{26}+a_8a_{10}^3x_{36}y_{20}y_{26}+a_{10}^2x_{36}y_{64},\\
y_{60}y_{76} &= a_4^3x_{48}x_{54}y_{22}+a_4^4a_8a_{10}x_{48}x_{54}-a_4^3a_8^5a_{10}x_{54}y_{20}+a_4^3a_8^3a_{10}^3x_{48}y_{22}-a_4^2a_8^4x_{54}y_{20}y_{22}\\
&\qquad -a_4^2a_8^7a_{10}^3y_{20}y_{22}+a_4^2a_8^6a_{10}^2y_{60}-a_4^2a_8^5a_{10}^3y_{58}+a_4^2a_8^4a_{10}^2y_{76}-a_4^2a_8a_{10}^3x_{48}y_{20}y_{22}\\
&\qquad +a_4^2a_{10}^2x_{48}y_{60}+a_4a_8^4a_{10}x_{36}x_{54}+a_4a_8a_{10}^4x_{36}x_{48}+a_8^3x_{36}x_{54}y_{22}\\
&\qquad +a_8^4a_{10}^2x_{36}y_{22}y_{26}+a_8^2a_{10}x_{36}x_{54}y_{20}-a_8^2a_{10}^2x_{36}y_{64}-a_{10}^3x_{36}x_{48}y_{22},\\
y_{60}y_{26} &= -a_4a_8x_{54}y_{20}-a_8^3a_{10}^2y_{20}y_{22}+a_8^2a_{10}y_{60}-a_8a_{10}^2y_{58}-a_{10}y_{76},
\end{aligned}
$$

$$\begin{aligned}
y_{60}y_{64} &= a_4^4a_{10}x_{48}x_{54} + a_4^3a_8^4a_{10}x_{54}y_{20} + a_4^3a_8^2a_{10}^3x_{48}y_{22} - a_4^2a_8x_{54}y_{58} \\
&\quad - a_4^2a_8^3x_{54}y_{20}y_{22} + a_4^2a_8^6a_{10}^3y_{20}y_{22} + a_4^2a_8^5a_{10}^2y_{60} + a_4^2a_8^4a_{10}^3y_{58} + a_4^2a_8a_{10}^2x_{48}y_{22}^2 \\
&\quad - a_4^2a_{10}^3x_{48}y_{20}y_{22} + a_4a_8^6a_{10}^4x_{36} - a_4a_8^3a_{10}x_{36}x_{54} \\
&\quad - a_4a_{10}^4x_{36}x_{48} + a_8^5a_{10}^3x_{36}y_{22} - a_8^4a_{10}^4x_{36}y_{20}, \\
y_{26}^3 &= a_8^3x_{54} - a_8^4a_{10}^2y_{26} - a_8^2a_{10}y_{26}^2 + a_{10}^3x_{48}, \\
y_{64}y_{26} &= -a_4a_8^4x_{54} + a_8^2x_{54}y_{20} - a_8^6a_{10}^2y_{22} + a_8^4a_{10}y_{22}y_{26} + a_8^2a_{10}y_{64} + a_{10}^2x_{48}y_{22}, \\
y_{64}^2 &= -a_4^2a_8^7a_{10}x_{54} + a_4^2a_8a_{10}x_{48}x_{54} - a_4a_8^6x_{54}y_{22} - a_4a_8^9a_{10}^3y_{22} + a_4a_8^5a_{10}x_{54}y_{20} \\
&\quad + a_4a_8^3a_{10}^3x_{48}y_{22} + a_8^2x_{54}y_{58} - a_8^4x_{54}y_{20}y_{22} - a_8^7a_{10}^3y_{20}y_{22} + a_8^5a_{10}^3y_{58} \\
&\quad - a_8^4a_{10}^2y_{76} - a_8^2a_{10}^2x_{48}y_{22}^2 - a_8a_{10}^3x_{48}y_{20}y_{22} - a_{10}^2x_{48}y_{60}, \\
y_{76}^2 &= -a_4^2x_{48}x_{54}y_{20}y_{22} - a_4^4a_8^3a_{10}x_{48}x_{54} - a_4^3a_8^2x_{48}x_{54}y_{22} + a_4^3a_8^7a_{10}x_{54}y_{20} \\
&\quad + a_4^3a_8^5a_{10}^3x_{48}y_{22} - a_4^3a_8a_{10}x_{48}x_{54}y_{20} - a_4^2a_8^4x_{54}y_{58} + a_4^2a_8^6x_{54}y_{20}y_{22} \\
&\quad - a_4^2a_8^9a_{10}^3y_{20}y_{22} - a_4^2a_8^8a_{10}^2y_{60} - a_4^2a_8^7a_{10}^3y_{58} + a_4^2a_8^5a_{10}x_{54}y_{20}^2 \\
&\quad - a_4^2a_8^4a_{10}^2x_{48}y_{22}^2 - a_4^2a_8a_{10}^3x_{48}y_{58} + a_4^2a_{10}^2x_{48}y_{76} - a_4a_8^9a_{10}^4x_{36} \\
&\quad - a_4a_8^3a_{10}^4x_{36}x_{48} + a_8^2x_{36}x_{54}y_{20}y_{26} - a_8^5x_{36}x_{54}y_{22} + a_8^8a_{10}^3x_{36}y_{22} \\
&\quad - a_8^7a_{10}^4x_{36}y_{20} - a_8^5a_{10}^3x_{36}y_{20}y_{26} + a_8^4a_{10}^2x_{36}y_{64} - a_8^2a_{10}^3x_{36}x_{48}y_{22} \\
&\quad + a_{10}^2x_{36}x_{48}y_{22}y_{26}, \\
y_{76}y_{26} &= a_4a_8^3x_{54}y_{20} - a_4a_8a_{10}^2x_{48}y_{22} + a_8x_{54}y_{20}^2 - a_8^5a_{10}^2y_{20}y_{22} \\
&\quad + a_8^4a_{10}y_{60} - a_8^3a_{10}^2y_{58} - a_8^2a_{10}y_{76} + a_{10}x_{48}y_{22}^2, \\
y_{64}y_{76} &= -a_4^3a_8^2a_{10}x_{48}x_{54} - a_4^2a_8x_{48}x_{54}y_{22} - a_4^2a_8^6a_{10}x_{54}y_{20} - a_4^2a_{10}x_{48}x_{54}y_{20} \\
&\quad - a_4a_8^3x_{54}y_{58} - a_4a_8^5x_{54}y_{20}y_{22} + a_4a_8^8a_{10}^3y_{20}y_{22} + a_4a_8^7a_{10}^2y_{60} \\
&\quad + a_4a_8^6a_{10}^3y_{58} - a_4a_8^5a_{10}^2y_{76} - a_4a_8^4a_{10}x_{54}y_{20}^2 - a_4a_8^3a_{10}^2x_{48}y_{22}^2 \\
&\quad + a_4a_8a_{10}^2x_{48}y_{60} - a_4a_{10}^3x_{48}y_{58} + a_8^3x_{36}x_{54}y_{26} + a_8^8a_{10}^4x_{36} \\
&\quad - a_8^6a_{10}^3x_{36}y_{26} + a_8^5a_{10}x_{36}x_{54} + a_8^5a_{10}^2y_{58}y_{22} + a_8^4a_{10}^2x_{36}y_{26}^2 \\
&\quad + a_8^2a_{10}^4x_{36}x_{48} + a_{10}^3x_{36}x_{48}y_{26}.
\end{aligned}$$

ii) $a_9^2,\ y_{21}^2,\ y_{25}^2,\ y_{27}^2,$

$a_9y_{21} + x_{26}a_4,\ a_9y_{25} + x_{26}a_8,\ a_9y_{27} + x_{26}a_{10},$

$y_{21}y_{25} + x_{26}y_{20},\ y_{21}y_{27} - x_{26}y_{22},\ y_{25}y_{27} - x_{26}y_{26},$

iii) $a_9\partial^2Q,\ y_{21}\partial^2Q,\ y_{25}\partial^2Q,\ y_{27}\partial^2Q,\ x_{26}\partial^2Q,$

$a_9a_4,\ a_9a_8,\ a_9a_{10},$

$y_{21}a_4,\ y_{25}a_8,\ y_{27}a_{10},$

$y_{21}a_8 + a_9y_{20} = y_{25}a_4 - a_9y_{20},$

$y_{21}a_{10} - a_9y_{22} = y_{27}a_4 + a_9y_{22},$

$y_{25}a_{10} - a_9y_{26} = y_{27}a_8 + a_9y_{26},$

$y_{21}y_{20},\ y_{25}y_{20},\ y_{21}y_{22},\ y_{27}y_{22},\ y_{25}y_{26},\ y_{27}y_{26},$

$y_{27}y_{20} - y_{25}y_{22} = y_{25}y_{22} + y_{21}y_{26} = -y_{27}y_{20} - y_{21}y_{26}.$

Here the definition of ∂ is given in (3.3), and the table of ∂^2-image is given in Lemma 4.2.

3. Proof of Theorem 2.1

First let us recall some results from [MS]. We consider the differential algebra $(\overline{V}, d)$ defined as follows:

$$\overline{V} = T(a_4, a_8, a_{10}, b_{12}, b_{16}, b_{18}, a_9, c_{17})/J,$$

where J is the ideal generated by

$$a_9 b_j - b_j a_9 + c_{17} a_{j-8} \quad \text{for} \quad j = 12, 16, 18$$

and

$$\alpha\beta - (-1)^{|\beta||\alpha|}\beta\alpha \qquad \text{for all the pairs } (\alpha, \beta) \text{ of the generators except } (a_9, b_j) \text{ with } j = 12, 16, 18 \text{ and } (a_9, c_{17});$$

the differential d is given over generators by

$$da_i = 0 \quad \text{for} \quad i = 4, 8, 9, 10, \quad dc_{17} = a_9^2, \quad db_j = -a_9 a_{j-8} \quad \text{for} \quad j = 12, 16, 18. \tag{3.1}$$

Then the differential algebra $(\overline{V}, d)$ is seen to be a resolution of $\mathbb{Z}_3$ over $H^*(E_6)$. (For a proof, see Theorem 5.7 and Corollary 5.8 in [MS].) So by the definition of Cotor we have

Theorem ([MS]) $\quad H^*(\overline{V}, d) = \mathrm{Cotor}_{H^*(E_6)}(\mathbb{Z}_3, \mathbb{Z}_3).$

We will use a spectral sequence to calculate $H^*(\overline{V}, d)$. To that end we need to introduce a filtration in $\overline{V}$. To begin with, we fix an additive basis of $\overline{V}$. The following lemma is straightforward from the definition of the ideal J.

Lemma 3.1. *We have an additive isomorphism*

$$\overline{V} \cong T(a_9, c_{17}) \otimes \mathbb{Z}_3[a_4, a_8, a_{10}, b_{12}, b_{16}, b_{18}].$$

Hence as a basis of $\overline{V}$ we can choose the following monomials:

$$a_9^{\ell_1} c_{17}^{m_1} \dots a_9^{\ell_p} c_{17}^{m_p} a_4^{i_1} a_8^{i_2} a_{10}^{i_3} b_{12}^{j_1} b_{16}^{j_2} b_{18}^{j_3}. \tag{3.2}$$

We define a weight by

$$\begin{aligned} w(a_i) &= 2 \qquad \text{for } i = 4, 8, 10; \\ w(b_j) &= 0 \qquad \text{for } j = 12, 16, 18; \\ w(a_9) &= 1, \quad w(c_{17}) = 2, \quad w(0) = \infty. \end{aligned}$$

For a monomial of the form (3.2), we define the weight by

$$w(\alpha_1 \dots \alpha_n) = w(\alpha_1) + \cdots + w(\alpha_n).$$

For an element $x = \sum \lambda_i x_i$, where x_i is a monomial given in (3.2) and $\lambda_i \in \mathbb{Z}_3$, we define that $w(x) = \inf_i w(x_i)$. Then we introduce a filtration in $\overline{V}$ by setting

$$F^p\overline{V} = \{x \in \overline{V} \mid w(x) \geq p\} \quad \text{for} \quad p \in \mathbb{N}.$$

It is easily seen that $\{F^p\overline{V}\}$ is a family of differential graded algebras giving a decreasing filtration. Hence from this filtration we can construct a spectral sequence $\{E_r, d_r\}$ such that $E_0 = \overline{V}$ and $E_\infty = H^*(\overline{V}, d)$, namely, converging to $\mathrm{Cotor}_{H^*(E_6)}(\mathbb{Z}_3, \mathbb{Z}_3)$.

It is convenient for later use to introduce here an auxiliary derivation ∂ on a subalgebra $\mathbb{Z}_3[a_4, a_8, a_{10}, b_{12}, b_{16}, b_{18}]$ of $\overline{V}$ defined as in (5.9) of [MS]:

$$\partial a_i = 0 \quad \text{for} \quad i = 4, 8, 10; \quad \partial b_j = -a_{j-8} \quad \text{for} \quad j = 12, 16, 18. \tag{3.3}$$

Since the only non trivial differential on the generators of E_0 is $d_0(c_{17}) = a_9^2$ by (3.1), we have that

$$E_1 = \mathbb{Z}_3[x_{26}] \otimes \mathbb{Z}_3[a_4, a_8, a_{10}] \otimes \Lambda(a_9) \otimes \mathbb{Z}_3[b_{12}, b_{16}, b_{18}],$$

where $x_{26} = a_9c_{17} - c_{17}a_9$. Since $d_1 = d_2 = 0$ for degree reasons, we have

$$E_1 = E_2 = E_3.$$

We remark here that $E_3 = E_1$ is a commutative algebra in the graded sense and also that d_3 is a derivative differential, i.e.,

$$d_3(xy) = d_3(x)y \pm xd_3(y), \qquad x, y \in E_3$$

such that

$$d_3(b_{12}) = -a_4a_9, \quad d_3(b_{16}) = -a_8a_9, \quad d_3(b_{18}) = -a_{10}a_9,$$

which are obtained by the differential formula (3.1).

In the next steps, we will use the following

Convention Let (M, d) be a differential module such that $M = M_1 \oplus M_2 \oplus N_1$ and that $d(M_1) \subset N_1$ and $d|_{M_2 \oplus N_1} = 0$. Then we denote $\mathrm{Ker}\ d \cap M_1$ by $\mathrm{Ker}\ d$ and $N_1/d(M_1)$ by $\mathrm{Coker}\ d$.

Under our convention, we have that

$$H(M) = \mathrm{Ker}\ d \oplus \mathrm{Coker}\ d \oplus M_2.$$

Moreover, for simplicity we will describe the differential d only on M_1. If not specifically denoted, we understand that $d(a) = 0$ for $a \in M_2 \oplus N_1$.

To calculate the E_4-term, we introduce another spectral sequence starting with the E_3-term. That is, we define a weight $\widetilde{w}$ by

$$\widetilde{w}(b_i) = 0 \quad \text{for } i = 12, 16, 18,\ \widetilde{w}(x_{26}) = \widetilde{w}(a_9) = \widetilde{w}(a_4) = 0,\ \widetilde{w}(a_8) = 1,\ \widetilde{w}(a_{10}) = 3. \tag{3.4}$$

By similarly introducing a filtration using the weight $\widetilde{w}$ as before, we obtain a new spectral sequence $\{\widetilde{E}_r, \widetilde{d}_r\}$ such that $\widetilde{E}_0 = E_3$ and $\widetilde{E}_\infty = E_4$.

At the $\widetilde{E}_0$-term we see that the relation $d_3(b_{12}) = -a_4a_9$ gives $\widetilde{d}_0(b_{12}) = -a_4a_9$, from which we obtain that

$$\widetilde{E}_1 = \Big(\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{1, a_9b_{12}^2\} \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9, a_9b_{12}\}\Big) \otimes \mathbb{Z}_3[b_{16}, b_{18}] \otimes \mathbb{Z}_3[b_{12}^3].$$

Since the relation $d_3(b_{16}) = -a_8a_9$ gives $\widetilde{d}_1(b_{16}) = -a_9a_8$, we have that

$$\begin{aligned}\widetilde{E}_2 = \Big(&\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}] \oplus \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{a_4b_{16}, a_4b_{16}^2\}\\ &\oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9, a_9b_{16}\} \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{16}^2\}\\ &\oplus \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{a_9b_{12}^2, a_9b_{12}^2b_{16}, a_9b_{12}^2b_{16}^2\}\\ &\oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{12}, a_9b_{12}b_{16}, a_9b_{12}b_{16}^2\}\Big) \otimes \mathbb{Z}_3[b_{18}] \otimes \mathbb{Z}_3[b_{12}^3, b_{16}^3].\end{aligned}$$

It follows from the relation $d(a_4b_{16}^2 + a_8b_{12}b_{16}) = -a_8^2a_9b_{12} + a_4a_8^2c_{17}$ giving $d_3(a_4b_{16}^2 + a_8b_{12}b_{16}) = -a_8^2a_9b_{12}$ in E_3 that $\widetilde{d}_2(a_4b_{16}^2) = -a_8^2a_9b_{12}$, which implies that

$$\begin{aligned}\widetilde{E}_3 = \Big(&\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{1, a_4b_{16}, a_4^2b_{16}^2\} \oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9, a_9b_{16}\}\\ &\oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{16}^2\} \oplus \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{a_9b_{12}^2, a_9b_{12}^2b_{16}, a_9b_{12}^2b_{16}^2\}\\ &\oplus \mathbb{Z}_3[a_{10}, x_{26}] \otimes \Delta(a_8)\{a_9b_{12}\} \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{12}b_{16}, a_9b_{12}b_{16}^2\}\Big)\\ &\otimes \mathbb{Z}_3[b_{12}^3, b_{16}^3] \otimes \mathbb{Z}_3[b_{18}].\end{aligned}$$

Using the expression $\mathbb{Z}_3[b_{18}] = \mathbb{Z}_3\{1, b_{18}, b_{18}^2\} \otimes \mathbb{Z}_3[b_{18}^3]$, the $\widetilde{E}_3$-term may be rewritten as

$$\begin{aligned}\widetilde{E}_3 = \Big(&\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{1, b_{18}, b_{18}^2, a_4b_{16}, a_4b_{16}b_{18}, a_4b_{16}b_{18}^2, a_4^2b_{16}^2, a_4^2b_{16}^2b_{18}, a_4^2b_{16}^2b_{18}^2\}\\ &\oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9, a_9b_{18}, a_9b_{18}^2, a_9b_{16}, a_9b_{16}b_{18}, a_9b_{16}b_{18}^2\}\\ &\oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{16}^2, a_9b_{16}^2b_{18}, a_9b_{16}^2b_{18}^2\} &&(4)\\ &\oplus \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{a_9b_{12}^2, a_9b_{12}^2b_{18}, a_9b_{12}^2b_{18}^2, a_9b_{12}^2b_{16}, a_9b_{12}^2b_{16}b_{18}, a_9b_{12}^2b_{16}b_{18}^2,\\ &\qquad a_9b_{12}^2b_{16}^2, a_9b_{12}^2b_{16}^2b_{18}, a_9b_{12}^2b_{16}^2b_{18}^2\} &&(5)\\ &\oplus \mathbb{Z}_3[a_{10}, x_{26}] \otimes \Delta(a_8)\{a_9b_{12}, a_9b_{12}b_{18}, a_9b_{12}b_{18}^2\}\\ &\oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{12}b_{16}, a_9b_{12}b_{16}b_{18}, a_9b_{12}b_{16}b_{18}^2,\\ &\qquad a_9b_{12}b_{16}^2, a_9b_{12}b_{16}^2b_{18}, a_9b_{12}b_{16}^2b_{18}^2\}\Big) &&(7)\\ &\otimes \mathbb{Z}_3[b_{12}^3, b_{16}^3, b_{18}^3],\end{aligned}$$

where (4), (5) and (7) will appear in Lemma 3.6. The only non trivial differential

$$\widetilde{d}_3 : \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{b_{18}, b_{18}^2\} \longrightarrow \mathbb{Z}_3[a_{10}, x_{26}]\{a_9, a_9b_{18}\}$$

is given by $\widetilde{d}_3(b_{18}) = -a_9a_{10}$ which is obtained from $d(b_{18}) = -a_9a_{10}$, and hence $\widetilde{d}_3(b_{18}^2) = a_9a_{10}b_{18}$. It is immediate to see that

$$\begin{aligned}\operatorname{Ker} \widetilde{d}_3 &= \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{a_4b_{18}, a_4b_{18}^2\} \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_8b_{18}, a_8b_{18}^2\}, &&\underline{1}'\\ \operatorname{Coker} \widetilde{d}_3 &= \mathbb{Z}_3[x_{26}]\{a_9, a_9b_{18}\}. &&\underline{2}'\end{aligned}$$

Thus we have that

$$\begin{aligned}
\widetilde{E}_4 = \Big(&\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{a_4b_{18},a_4b_{18}^2\}\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_8b_{18},a_8b_{18}^2\}\\
&\oplus\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{1,a_4b_{16},a_4b_{16}b_{18},a_4b_{16}b_{18}^2,a_4^2b_{16}^2,a_4^2b_{16}^2b_{18},a_4^2b_{16}^2b_{18}^2\}\\
&\oplus\mathbb{Z}_3[x_{26}]\{a_9,a_9b_{18}\}\\
&\oplus\mathbb{Z}_3[a_{10},x_{26}]\{a_9b_{18}^2,a_9b_{16},a_9b_{16}b_{18},a_9b_{16}b_{18}^2\}\\
&\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_9b_{16}^2,a_9b_{16}^2b_{18},a_9b_{16}^2b_{18}^2\}\\
&\oplus\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{a_9b_{12}^2,a_9b_{12}^2b_{18},a_9b_{12}^2b_{18}^2,a_9b_{12}^2b_{16},a_9b_{12}^2b_{16}b_{18},a_9b_{12}^2b_{16}b_{18}^2,\\
&\qquad\qquad a_9b_{12}^2b_{16}^2,a_9b_{12}^2b_{16}^2b_{18},a_9b_{12}^2b_{16}^2b_{18}^2\}\\
&\oplus\mathbb{Z}_3[a_{10},x_{26}]\otimes\Delta(a_8)\{a_9b_{12},a_9b_{12}b_{18},a_9b_{12}b_{18}^2\}\\
&\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_9b_{12}b_{16},a_9b_{12}b_{16}b_{18},a_9b_{12}b_{16}b_{18}^2,\\
&\qquad\qquad a_9b_{12}b_{16}^2,a_9b_{12}b_{16}^2b_{18},a_9b_{12}b_{16}^2b_{18}^2\}\Big)\\
&\otimes\mathbb{Z}_3[b_{12}^3,b_{16}^3,b_{18}^3].
\end{aligned}$$

The relation $d_3(a_4b_{16}b_{18}-a_8b_{12}b_{18}-a_{10}b_{12}b_{16})=-a_8a_9a_{10}b_{12}$ in E_3 implies that the only non trivial differential

$$\widetilde{d}_4:\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{a_4b_{16}b_{18},a_4b_{16}b_{18}^2\}\to\mathbb{Z}_3[a_{10},x_{26}]\otimes\Delta(a_8)\{a_9b_{12},a_9b_{12}b_{18}\}$$

is given by $\widetilde{d}_4(a_4b_{16}b_{18})=-a_8a_9a_{10}b_{12}$, and hence $\widetilde{d}_4(a_4b_{16}b_{18}^2)=a_8a_9a_{10}b_{12}b_{18}$. So we have that

$$\begin{aligned}
\mathrm{Ker}\,\widetilde{d}_4 &= \mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{a_4^2b_{16}b_{18},a_4^2b_{16}b_{18}^2\}\\
&\qquad\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_4a_8b_{16}b_{18},a_4a_8b_{16}b_{18}^2\}, && \underline{3}'\\
\mathrm{Coker}\,\widetilde{d}_4 &= \mathbb{Z}_3[a_{10},x_{26}]\{a_9b_{12},a_9b_{12}b_{18}\}\oplus\mathbb{Z}_3[x_{26}]\{a_9a_8b_{12},a_9a_8b_{12}b_{18}\}. && \underline{4}'
\end{aligned}$$

Thus we have that

$$\begin{aligned}
\widetilde{E}_5 = \Big(&\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{a_4b_{18},a_4b_{18}^2\}\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_8b_{18},a_8b_{18}^2\}\\
&\oplus\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{1,a_4b_{16},a_4^2b_{16}^2,a_4^2b_{16}^2b_{18},a_4^2b_{16}^2b_{18}^2\}\\
&\oplus\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{a_4^2b_{16}b_{18},a_4^2b_{16}b_{18}^2\}\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_4a_8b_{16}b_{18},a_4a_8b_{16}b_{18}^2\}\\
&\oplus\mathbb{Z}_3[x_{26}]\{a_9,a_9b_{18}\}\oplus\mathbb{Z}_3[a_{10},x_{26}]\{a_9b_{18}^2,a_9b_{16},a_9b_{16}b_{18},a_9b_{16}b_{18}^2\}\\
&\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_9b_{16}^2,a_9b_{16}^2b_{18},a_9b_{16}^2b_{18}^2\}\\
&\oplus\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{a_9b_{12}^2,a_9b_{12}^2b_{18},a_9b_{12}^2b_{18}^2,a_9b_{12}^2b_{16},a_9b_{12}^2b_{16}b_{18},a_9b_{12}^2b_{16}b_{18}^2,\\
&\qquad\qquad a_9b_{12}^2b_{16}^2,a_9b_{12}^2b_{16}^2b_{18},a_9b_{12}^2b_{16}^2b_{18}^2\}\\
&\oplus\mathbb{Z}_3[a_{10},x_{26}]\{a_9b_{12},a_9b_{12}b_{18}\}\oplus\mathbb{Z}_3[x_{26}]\{a_9a_8b_{12},a_9a_8b_{12}b_{18}\}\\
&\oplus\mathbb{Z}_3[a_{10},x_{26}]\otimes\Delta(a_8)\{a_9b_{12}b_{18}^2\}\\
&\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_9b_{12}b_{16},a_9b_{12}b_{16}b_{18},a_9b_{12}b_{16}b_{18}^2,\\
&\qquad\qquad a_9b_{12}b_{16}^2,a_9b_{12}b_{16}^2b_{18},a_9b_{12}b_{16}^2b_{18}^2\}\Big)\\
&\otimes\mathbb{Z}_3[b_{12}^3,b_{16}^3,b_{18}^3].
\end{aligned}$$

The relation $d_3(a_8b_{18}^2+a_{10}b_{16}b_{18})=-a_{10}^2a_9b_{16}$ in E_3 implies that the only non trivial differential

$$\widetilde{d_5}:\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_8b_{18}^2\}\to\mathbb{Z}_3[a_{10},x_{26}]\{a_9b_{16}\}$$

is given by $\widetilde{d_5}(a_8b_{18}^2)=-a_{10}^2a_9b_{16}$. Hence we obtain that

$$\begin{aligned}\operatorname{Ker}\widetilde{d_5}&=\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_8^2b_{18}^2\}, && \underline{5}'\\ \operatorname{Coker}\widetilde{d_5}&=\Delta(a_{10})\otimes\mathbb{Z}_3[x_{26}]\{a_9b_{16}\}. && \underline{6}'\end{aligned}$$

Thus we have that

$$\begin{aligned}\widetilde{E}_6=\Big(&\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{a_4b_{18},a_4b_{18}^2\}\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_8b_{18},a_8^2b_{18}^2\}\\ &\oplus\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{1,a_4b_{16},a_4^2b_{16}^2,a_4^2b_{16}^2b_{18},a_4^2b_{16}^2b_{18}^2,a_4^2b_{16}b_{18},a_4^2b_{16}b_{18}^2\}\\ &\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_4a_8b_{16}b_{18},a_4a_8b_{16}b_{18}^2\}\\ &\oplus\mathbb{Z}_3[x_{26}]\{a_9,a_9b_{18},a_9a_8b_{12},a_9a_8b_{12}b_{18}\}\\ &\oplus\mathbb{Z}_3[a_{10},x_{26}]\{a_9b_{18}^2,a_9b_{16}b_{18},a_9b_{16}b_{18}^2,a_9b_{12},a_9b_{12}b_{18}\}\\ &\oplus\mathbb{Z}_3[x_{26}]\otimes\Delta(a_{10})\{a_9b_{16}\}\\ &\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_9b_{16}^2,a_9b_{16}^2b_{18},a_9b_{16}^2b_{18}^2\}\\ &\oplus\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{a_9b_{12}^2,a_9b_{12}^2b_{18},a_9b_{12}^2b_{18}^2,a_9b_{12}^2b_{16},a_9b_{12}^2b_{16}b_{18},a_9b_{12}^2b_{16}b_{18}^2,\\ &\qquad\qquad a_9b_{12}^2b_{16}^2,a_9b_{12}^2b_{16}^2b_{18},a_9b_{12}^2b_{16}^2b_{18}^2\}\\ &\oplus\mathbb{Z}_3[a_{10},x_{26}]\otimes\Delta(a_8)\{a_9b_{12}b_{18}^2\}\\ &\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_9b_{12}b_{16},a_9b_{12}b_{16}b_{18},a_9b_{12}b_{16}b_{18}^2,\\ &\qquad\qquad a_9b_{12}b_{16}^2,a_9b_{12}b_{16}^2b_{18},a_9b_{12}b_{16}^2b_{18}^2\}\Big)\\ &\otimes\mathbb{Z}_3[b_{12}^3,b_{16}^3,b_{18}^3].\end{aligned}$$

Similarly the relation $d_3(a_4b_{18}^2)=a_9a_{10}^2b_{12}$ in E_3 implies that the only non trivial differential

$$\widetilde{d_6}:\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{a_4b_{18}^2\}\to\mathbb{Z}_3[a_{10},x_{26}]\{a_9b_{12}\}$$

is given by $\widetilde{d_6}(a_4b_{18}^2)=a_9a_{10}^2b_{12}$, from which we have that

$$\begin{aligned}\operatorname{Ker}\widetilde{d_6}&=\mathbb{Z}_3[a_4,a_8,a_{10},x_{26}]\{a_4^2b_{18}^2\}\oplus\mathbb{Z}_3[a_8,a_{10},x_{26}]\{a_4a_8b_{18}^2\}, && \underline{7}'\\ \operatorname{Coker}\widetilde{d_6}&=\mathbb{Z}_3[x_{26}]\otimes\Delta(a_{10})\{a_9b_{12}\}. && \underline{8}'\end{aligned}$$

Thus we have that

$$\begin{aligned}\widetilde{E}_7 = \Big(&\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{a_4b_{18}, a_4^2b_{18}^2\} \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_8b_{18}, a_8b_{18}^2, a_4a_8b_{18}^2\}\\
&\oplus \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{1, a_4b_{16}, a_4^2b_{16}^2, a_4^2b_{16}^2b_{18}, a_4^2b_{16}^2b_{18}^2, a_4^2b_{16}b_{18}, a_4^2b_{16}b_{18}^2\}\\
&\oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_4a_8b_{16}b_{18}, a_4a_8b_{16}b_{18}^2\}\\
&\oplus \mathbb{Z}_3[x_{26}]\{a_9, a_9b_{18}, a_9a_8b_{12}, a_9a_8b_{12}b_{18}\}\\
&\oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{18}^2, a_9b_{16}b_{18}, a_9b_{16}b_{18}^2, a_9b_{12}b_{18}\}\\
&\oplus \mathbb{Z}_3[x_{26}] \otimes \Delta(a_{10})\{a_9b_{12}, a_9b_{16}\}\\
&\oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{16}^2, a_9b_{16}^2b_{18}, a_9b_{16}^2b_{18}^2\}\\
&\oplus \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{a_9b_{12}^2, a_9b_{12}^2b_{18}, a_9b_{12}^2b_{18}^2, a_9b_{12}^2b_{16}, a_9b_{12}^2b_{16}b_{18}, a_9b_{12}^2b_{16}b_{18}^2,\\
&\qquad\qquad a_9b_{12}^2b_{16}^2, a_9b_{12}^2b_{16}^2b_{18}, a_9b_{12}^2b_{16}^2b_{18}^2\}\\
&\oplus \mathbb{Z}_3[a_{10}, x_{26}] \otimes \Delta(a_8)\{a_9b_{12}b_{18}^2\}\\
&\oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{12}b_{16}, a_9b_{12}b_{16}b_{18}, a_9b_{12}b_{16}b_{18}^2,\\
&\qquad\qquad a_9b_{12}b_{16}^2, a_9b_{12}b_{16}^2b_{18}, a_9b_{12}b_{16}^2b_{18}^2\}\Big)\\
\otimes\, &\mathbb{Z}_3[b_{12}^3, b_{16}^3, b_{18}^3].\end{aligned}$$

Then for degree reasons we see that $\widetilde{d}_r = 0$ for $r \geq 7$ so that $\widetilde{E}_7 = \widetilde{E}_\infty$, which gives the E_4-term of the original spectral sequence.

Now we can summarize our calculations from (4) to (7) and from $\underline{1}'$ to $\underline{8}'$ as follows; for the kernel parts, we have

$$\begin{aligned}&\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{1, a_4b_{16}, a_4^2b_{16}^2, a_4b_{18}, a_4^2b_{18}^2, a_4^2b_{16}b_{18}, a_4^2b_{16}^2b_{18}, a_4^2b_{16}b_{18}^2, a_4^2b_{16}^2b_{18}^2\}\\
&\qquad \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_8b_{16}, a_8^2b_{16}^2, a_4a_8b_{16}b_{18}, a_4a_8b_{18}^2, a_4a_8b_{16}b_{18}^2\}; \end{aligned} \tag{1}$$

for the cokernel parts, we have

$$\begin{aligned}&\mathbb{Z}_3[x_{26}]\{a_9, a_9b_{12}, a_9b_{16}, a_9b_{18}, a_9a_8b_{12}, a_9a_{10}b_{12}, a_9a_{10}b_{16}, a_9a_8b_{12}b_{18}\}\\
&\qquad \oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{12}b_{18}\}.\end{aligned} \tag{2}$$

Hence we can write

$$\begin{aligned}\text{(3.5)}\quad \widetilde{E}_7 = \widetilde{E}_\infty = (1) \oplus (2) \oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{12}b_{18}\} \oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{16}b_{18}, a_9b_{16}b_{18}^2, a_9b_{18}^2\}\\
\oplus (4) \oplus (5) \oplus \mathbb{Z}_3[a_{10}, x_{26}] \otimes \Delta(a_8)\{a_9b_{12}b_{18}^2\} \oplus (7).\end{aligned}$$

Then one can verify that the following elements are chosen to be generators of the filtered algebra E_0:

$$\begin{array}{llll}\overline{y}_{20} = -a_4b_{16}, & \overline{y}_{22} = a_4b_{18}, & \overline{y}_{26} = a_8b_{18}, & \\
\overline{y}_{58} = -a_4^2b_{16}^2b_{18}, & \overline{y}_{60} = -a_4^2b_{16}b_{18}^2, & \overline{y}_{64} = -a_8^2b_{12}b_{18}^2, & \overline{y}_{76} = -a_4^2b_{16}^2b_{18}^2,\\
\overline{y}_{21} = a_9b_{12}, & \overline{y}_{25} = a_9b_{16}, & \overline{y}_{27} = a_9b_{18}. & \end{array}$$

Numbering the term $\mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{12}b_{18}, a_9b_{16}b_{18}, a_9b_{16}b_{18}^2, a_9b_{18}^2\}$ by (3) and the term $\mathbb{Z}_3[a_{10}, x_{26}] \otimes \Delta(a_8)\{a_9b_{12}b_{18}^2\}$ by (6), we have proved the following

Lemma 3.2. *We have that*

$$
\begin{aligned}
E_4 = \Big(& \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{1, y_{20}, y_{20}^2, y_{22}, y_{22}^2, y_{20}y_{22}, y_{58}, y_{60}, y_{76}\} \\
& \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{y_{26}, y_{26}^2, y_{20}y_{26}, y_{22}y_{26}, y_{64}\} && (1) \\
& \oplus \mathbb{Z}_3[x_{26}]\{a_9, y_{21}, y_{25}, y_{27}, y_{21}a_8, y_{21}a_{10}, y_{25}a_{10}, y_{21}y_{26}\} && (2) \\
& \oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{16}b_{18}, a_9b_{16}b_{18}^2, a_9b_{12}b_{18}, a_9b_{18}^2\} && (3) \\
& \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{16}^2, a_9b_{16}^2b_{18}, a_9b_{16}^2b_{18}^2\} && (4) \\
& \oplus \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]\{a_9b_{12}^2, a_9b_{12}^2b_{18}, a_9b_{12}^2b_{18}^2, a_9b_{12}^2b_{16}, a_9b_{12}^2b_{16}b_{18}, \\
& \qquad a_9b_{12}^2b_{16}b_{18}^2, a_9b_{12}^2b_{16}^2, a_9b_{12}^2b_{16}^2b_{18}, a_9b_{12}^2b_{16}^2b_{18}^2\} && (5) \\
& \oplus \mathbb{Z}_3[a_{10}, x_{26}] \otimes \Delta(a_8)\{a_9b_{12}b_{18}^2\} && (6) \\
& \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{12}b_{16}, a_9b_{12}b_{16}b_{18}, a_9b_{12}b_{16}b_{18}^2, \\
& \qquad a_9b_{12}b_{16}^2, a_9b_{12}b_{16}^2b_{18}, a_9b_{12}b_{16}^2b_{18}^2\}\Big) && (7) \\
& \otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}],
\end{aligned}
$$

where the cocycles y_i and x_i are defined as follows:

$$
\begin{array}{llll}
y_{20} = a_8b_{12} - a_4b_{16}, & y_{22} = a_4b_{18} - a_{10}b_{12}, & y_{26} = a_8b_{18} - a_{10}b_{16}, & \\
y_{58} = \partial^2(b_{12}^2b_{16}^2b_{18}), & y_{60} = \partial^2(b_{12}^2b_{16}b_{18}^2), & y_{64} = \partial^2(b_{12}b_{16}^2b_{18}^2), & y_{76} = \partial^2(b_{12}^2b_{16}^2b_{18}^2), \\
y_{21} = a_9b_{12} - c_{17}a_4, & y_{25} = a_9b_{16} - c_{17}a_8, & y_{27} = a_9b_{18} - c_{17}a_{10}, & \\
x_{36} = b_{12}^3, & x_{48} = b_{16}^3, & x_{54} = b_{18}^3. &
\end{array}
$$

As for the auxiliary derivation ∂ defined in (3.3) we have the following formula:

Lemma 3.3. *Let Q be an element of $\mathbb{Z}_3[a_4, a_8, a_{10}, b_{12}, b_{16}, b_{18}]$. Then we have*

$$x_{26}\partial^2(-Q) = d(a_9Q + c_{17}\partial Q). \tag{3.6}$$

Proof. Using a formula $b_i^n a_9 = a_9b_i^n + nc_{17}a_{i-8}b_i^{n-2}$, one can show by a tedious computation that

$$d(a_9Q) = -a_9^2\partial Q - a_9c_{17}\partial^2 Q \quad \text{and} \quad d(c_{17}\partial Q) = a_9^2\partial Q + c_{17}a_9\partial^2 Q,$$

from which the lemma follows. □

Looking at the expression of Lemma 3.2, we see that all the possible non permanent cocycles appear in the terms from (3) to (7). Consider an element

$$-x_{26}\partial^2 Q = d(a_9Q + c_{17}\partial Q)$$

in (3.6), where $Q = f(a_4, a_8, a_{10})b_{12}^i b_{16}^j b_{18}^k \in \mathbb{Z}_3[a_4, a_8, a_{10}, b_{12}, b_{16}, b_{18}]$. Then we have

$$w(a_9Q) = w(a_9) + w(Q) = 1 + w(f).$$

We also have

$$w(\partial Q) \geq 2 + w(f) \quad \text{and} \quad w(\partial^2 Q) \geq 4 + w(f),$$

since ∂Q and $\partial^2 Q$ are of the form

$$\sum a_\mu f(a_4, a_8, a_{10})b_{12}^k b_{16}^\ell b_{18}^m \quad \text{and} \quad \sum a_\mu a_\nu f(a_4, a_8, a_{10})b_{12}^p b_{16}^q b_{18}^r$$

respectively. Thus we have

$$\begin{aligned}&w(a_9Q + c_{17}\partial Q) = 1 + w(f);\\ &w(-x_{26}\partial^2 Q) = w(x_{26}) + w(\partial^2 Q) \geq 3 + 4 + w(f) = 6 + w(a_9Q + c_{17}\partial Q).\end{aligned}$$

So we have

$$w(-x_{26}\partial^2 Q) - w(a_9Q + c_{17}\partial Q) \geq 6.$$

Therefore by the formula (3.6) we see that the differential d increases the filtration degree defined in the paragraph below (3.2) by more than 5. This means that $d_4 = 0$ and $d_5 = 0$, whence $E_4 = E_5 = E_6$.

Now it follows from Lemma 3.3 that

$$d_6(a_9Q) = -x_{26}\partial^2 Q.$$

To calculate the E_7-term, it is more convenient to regard the $\widetilde{E}_7$-term (or equivalently the $E_4 = E_6$-term) expressed in Lemma 3.2 as being equipped with the differential d_6. Then the filtration of E_6 defined similarly by (3.4)

$$(3.4)' \qquad \begin{aligned}&\widetilde{w}'(b_i) = 0 \quad \text{for} \quad i = 12, 16, 18,\\ &\widetilde{w}'(x_{26}) = \widetilde{w}'(a_9) = \widetilde{w}'(a_4) = 0, \quad \widetilde{w}'(a_8) = 1, \quad \widetilde{w}'(a_{10}) = 3\end{aligned}$$

induces a spectral sequence $\{\widetilde{E}'_r, \widetilde{d}'_r\}$ such that $\widetilde{E}'_0 = E_6$ and $\widetilde{E}'_\infty = E_7$. Then we see that the differential $\widetilde{d}'_0$ is non trivial only on (5) which is mapped to (1), where we have by the above formula and (3.3) that

$$\begin{aligned}&\widetilde{d}'_0(a_9b_{12}^2) = -x_{26}a_4^2, && \widetilde{d}'_0(a_9b_{12}^2b_{18}) = -x_{26}a_4y_{22}, && \widetilde{d}'_0(a_9b_{12}^2b_{18}^2) = -x_{26}y_{22}^2,\\ &\widetilde{d}'_0(a_9b_{12}^2b_{16}) = -x_{26}a_4y_{20}, && \widetilde{d}'_0(a_9b_{12}^2b_{16}b_{18}) = -x_{26}y_{20}y_{22}, && \widetilde{d}'_0(a_9b_{12}^2b_{16}b_{18}^2) = -x_{26}y_{60},\\ &\widetilde{d}'_0(a_9b_{12}^2b_{16}^2) = -x_{26}y_{20}^2, && \widetilde{d}'_0(a_9b_{12}^2b_{16}^2b_{18}) = -x_{26}y_{58}, && \widetilde{d}'_0(a_9b_{12}^2b_{16}^2b_{18}^2) = -x_{26}y_{76}.\end{aligned}$$

It follows from these relations that

$$\begin{aligned}\text{Ker } \widetilde{d}'_0 &= 0,\\ \text{Coker } \widetilde{d}'_0 &= \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/(x_{26}a_4^2) \oplus \Big(\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/(x_{26}a_4)\Big)\{y_{20}, y_{22}\}\\ &\qquad \oplus \mathbb{Z}_3[a_4, a_8, a_{10}]\{y_{20}^2, y_{22}^2, y_{20}y_{22}, y_{58}, y_{60}, y_{76}\}.\end{aligned}$$

Thus we obtain that

$$\widetilde{E}'_1 = \text{Coker } \widetilde{d}'_0 \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{y_{26}, y_{26}^2, y_{20}y_{26}, y_{22}y_{26}, y_{64}\} \oplus (2) \oplus (3) \oplus (4) \oplus (6) \oplus (7).$$

Now the differential $\widetilde{d}'_1$ is non trivial only on (7) which is mapped to the following

$$\text{Coker } \widetilde{d}'_0 \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{y_{26}, y_{26}^2, y_{20}y_{26}, y_{22}y_{26}, y_{64}\}.$$

Here to determine $\widetilde{d}'_1(a_9b_{12}b_{16}b_{18})$, we use the formula

$$a_4y_{26} + a_8y_{22} = -a_8y_{22} + a_{10}y_{20} = -a_{10}y_{20} - a_4y_{26} = \partial^2(b_{12}b_{16}b_{18}),$$

from which we see that $-a_8y_{22} + a_{10}y_{20} = -a_8y_{22}$ on $\widetilde{E}'_1$, since $-a_8y_{22}$ and $a_{10}y_{20}$ are of filtrations 1 and 2 respectively, and hence we have $\widetilde{d}'_1(a_9b_{12}b_{16}b_{18}) = a_8x_{26}y_{22}$. For the other elements of (7), the calculation of $\widetilde{d}'_1$ is straightforward from (3.6), and the results are as follows:

$$\begin{aligned}&\widetilde{d}'_1(a_9b_{12}b_{16}) = -x_{26}a_4a_8, \quad \widetilde{d}'_1(a_9b_{12}b_{16}b_{18}^2) = -x_{26}y_{22}y_{26},\\ &\widetilde{d}'_1(a_9b_{12}b_{16}^2) = -x_{26}a_8y_{20}, \quad \widetilde{d}'_1(a_9b_{12}b_{16}^2b_{18}) = -x_{26}y_{20}y_{26}, \quad \widetilde{d}'_2(a_9b_{12}b_{16}^2b_{18}^2) = -x_{26}y_{64}.\end{aligned}$$

Hence we have that

$$\begin{aligned}\operatorname{Ker} \widetilde{d}'_1 &= 0,\\ \operatorname{Coker} \widetilde{d}'_1 &= \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/(x_{26}a_4^2, x_{26}a_4a_8)\\ &\quad \oplus \Big(\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/(x_{26}a_4, x_{26}a_8)\Big)\{y_{20}, y_{22}\}\\ &\quad \oplus \mathbb{Z}_3[a_4, a_8, a_{10}]\{y_{20}^2, y_{22}^2, y_{20}y_{22}, y_{58}, y_{60}, y_{76}\} \qquad \underline{1}''\\ &\quad \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{y_{26}, y_{26}^2, y_{64}\} \oplus \mathbb{Z}_3[a_8, a_{10}]\{y_{20}y_{26}, y_{22}y_{26}\}.\end{aligned}$$

Thus we obtain that

$$\widetilde{E}'_2 = \operatorname{Coker} \widetilde{d}'_1 \oplus (2) \oplus (3) \oplus (4) \oplus (6).$$

For degree reasons we see that the elements $a_9b_{16}b_{18}$, $a_9b_{16}b_{18}^2$, $a_9b_{18}^2$ in (3) are $\widetilde{d}'_2$-cycles and so they survive to $\widetilde{E}'_3$.

Observe that the differential $\widetilde{d}'_2$ is non trivial only on (4). The differential $\widetilde{d}'_2$ maps (4) to the following summands

$$\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/(x_{26}a_4^2, x_{26}a_4a_8) \oplus \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{y_{26}, y_{26}^2\}.$$

It follows from (3.6) that

$$\begin{aligned}&\widetilde{d}'_2(a_9b_{16}^2) = -x_{26}a_8^2, \quad \widetilde{d}'_2(a_9b_{16}^2b_{18}) = -x_{26}a_8y_{26}, \quad \widetilde{d}'_2(a_9b_{16}^2b_{18}^2) = -x_{26}y_{26}^2,\\ &\widetilde{d}'_2(a_9b_{12}b_{16}^2b_{18}^2) = -x_{26}y_{64}.\end{aligned}$$

First we calculate the differential

$$\widetilde{d}'_2 : \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{16}^2\} \to \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/(x_{26}a_4^2, x_{26}a_4a_8),$$

where we have that

$$\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4^2, a_4a_8) \cong \mathbb{Z}_3[a_4, a_8, a_{10}] \oplus \mathbb{Z}_3[a_4, a_8]/(a_4^2, a_4a_8) \otimes \mathbb{Z}_3[a_{10}] \otimes \mathbb{Z}_3[x_{26}]^+.$$

We see that

$$\begin{aligned}\operatorname{Ker} \widetilde{d}'_2 \mid \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{16}^2\} &= 0,\\ \operatorname{Coker} \widetilde{d}'_2 &\cong \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4^2, a_4a_8, a_8^2)\\ &\cong \mathbb{Z}_3[a_4, a_8, a_{10}] \oplus \mathbb{Z}_3[a_4, a_8]/(a_4^2, a_4a_8, a_8^2) \otimes \mathbb{Z}_3[a_{10}] \otimes \mathbb{Z}_3[x_{26}]^+. \qquad (3.7)\end{aligned}$$

Also we easily see that

$$\begin{aligned}\operatorname{Ker} \widetilde{d}'_2 \mid \mathbb{Z}_3[a_8, a_{10}, x_{26}]\{a_9b_{16}^2b_{18}, a_9b_{12}b_{16}^2b_{18}^2\} \oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{16}^2b_{18}^2\} &= 0,\\ \operatorname{Coker} \widetilde{d}'_2 &= \Big(\mathbb{Z}_3[a_8, a_{10}, x_{26}]/x_{26}a_8\Big)\{y_{26}\} \oplus \mathbb{Z}_3[a_8, a_{10}]\{y_{26}^2, y_{64}\}.\end{aligned}$$

When we set

$$\begin{aligned} F = \Big(& \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4^2, a_4a_8, a_8^2) \\ & \oplus \Big(\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/(x_{26}a_4, x_{26}a_8)\Big)\{y_{20}, y_{22}\} \\ & \oplus \mathbb{Z}_3[a_4, a_8, a_{10}]\{y_{20}^2, y_{22}^2, y_{20}y_{22}, y_{58}, y_{60}, y_{76}\} \\ & \oplus \Big(\mathbb{Z}_3[a_8, a_{10}, x_{26}]/x_{26}a_8\Big)\{y_{26}\} \\ & \oplus \mathbb{Z}_3[a_8, a_{10}]\{y_{20}y_{26}, y_{22}y_{26}, y_{26}^2, y_{64}\}\Big) \otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}], \end{aligned}$$

we obtain

$$\widetilde{E}_3' = F \oplus (2) \oplus (3) \oplus (6).$$

We observe that the differential $\widetilde{d}_3'$ is non trivial only on the summand

$$\mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{12}b_{18}\} \oplus \mathbb{Z}_3[a_{10}, x_{26}] \otimes \Delta(a_8)\{a_9b_{12}b_{18}^2\}.$$

It follows from (3.6) that

$$\widetilde{d}_3'(a_9b_{12}b_{18}) = -x_{26}a_4a_{10}, \quad \widetilde{d}_3'(a_9b_{12}b_{18}^2) = -x_{26}a_{10}y_{22}.$$

First we calculate

$$\widetilde{d}_3' : \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{12}b_{18}\} \to \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4^2, a_4a_8, a_8^2).$$

Using (3.7), we obtain that

$$\begin{aligned} \operatorname{Ker} \widetilde{d}_3' \mid \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{12}b_{18}\} &= 0, \\ \operatorname{Coker} \widetilde{d}_3' &\cong \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4^2, a_4a_8, a_4a_{10}, a_8^2) \\ &\cong \mathbb{Z}_3[a_4, a_8, a_{10}] \oplus \mathbb{Z}_3[a_4, a_8, a_{10}]/(a_4^2, a_4a_8, a_4a_{10}, a_8^2) \otimes \mathbb{Z}_3[x_{26}]^+. \end{aligned} \tag{3.8}$$

By a similar way, we see that

$$\begin{aligned} \operatorname{Ker} \widetilde{d}_3' \mid \mathbb{Z}_3[a_{10}, x_{26}] \otimes \Delta(a_8)\{a_9b_{12}b_{18}^2\} &= \mathbb{Z}_3[a_{10}, x_{26}]\{a_9a_8b_{12}b_{18}^2\}, \\ \operatorname{Coker} \widetilde{d}_3' \mid \mathbb{Z}_3[a_{10}, x_{26}] \otimes \Delta(a_8)\{a_9b_{12}b_{18}^2\} &= \mathbb{Z}_3[a_4, a_8, a_{10}]\{y_{22}\} \oplus \mathbb{Z}_3[x_{26}]^+\{y_{22}\}. \end{aligned}$$

Thus we obtain that

$$\begin{aligned} \widetilde{E}_4' = \Big(& \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4^2, a_4a_8, a_4a_{10}, a_8^2) \\ & \oplus \Big(\mathbb{Z}_3[a_8, a_{10}, x_{26}]/(x_{26}a_8)\Big)\{y_{26}\} \oplus \mathbb{Z}_3[a_8, a_{10}]\{y_{26}^2\} \\ & \oplus \Big(\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4, a_8)\Big)\{y_{20}\} \\ & \oplus \mathbb{Z}_3[a_4, a_8, a_{10}]\{y_{22}\} \oplus \mathbb{Z}_3[x_{26}]^+\{y_{22}\} \\ & \oplus \mathbb{Z}_3[a_4, a_8, a_{10}]\{y_{20}^2, y_{22}^2, y_{20}y_{22}, y_{58}, y_{60}, y_{76}\} \\ & \oplus \mathbb{Z}_3[a_8, a_{10}]\{y_{20}y_{26}, y_{22}y_{26}, y_{64}\} \\ & \oplus \mathbb{Z}_3[x_{26}]\{a_9, y_{21}, y_{25}, y_{27}, y_{21}a_8, y_{21}a_{10}, y_{25}a_{10}, y_{21}y_{26}\} \\ & \oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{16}b_{18}, a_9b_{16}b_{18}^2, a_9b_{18}^2\} \\ & \oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9a_8b_{12}b_{18}^2\}\Big) \\ & \otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}]. \end{aligned}$$

The following are the only non trivial differentials on $\widetilde{E}'_4$:

$$\begin{aligned}\widetilde{d}'_4 &: \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{16}b_{18}\} \to \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4^2, a_4a_8, a_4a_{10}, a_8^2),\\ \widetilde{d}'_4 &: \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{16}b_{18}^2\} \to \Big(\mathbb{Z}_3[a_8, a_{10}, x_{26}]/x_{26}a_8\Big)\{y_{26}\},\end{aligned}$$

which are given by

$$\widetilde{d}'_4(a_9b_{16}b_{18}) = -x_{26}a_8a_{10}, \quad \widetilde{d}'_4(a_9b_{16}b_{18}^2) = -x_{26}a_{10}y_{26}$$

respectively. Hence we have respectively that

$$\begin{aligned}\mathrm{Ker}\ \widetilde{d}'_4 \mid \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{16}b_{18}\} &= 0,\\ \mathrm{Coker}\ \widetilde{d}'_4 \mid \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{16}b_{18}\} &\\ = \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}&(a_4^2, a_4a_8, a_4a_{10}, a_8^2, a_8a_{10})\\ = \mathbb{Z}_3[a_4, a_8, a_{10}] \oplus \mathbb{Z}_3&[a_4, a_8, a_{10}]/(a_4^2, a_4a_8, a_4a_{10}, a_8^2, a_8a_{10}) \otimes \mathbb{Z}_3[x_{26}]^+\end{aligned}$$

$$= \mathbb{Z}_3[a_4, a_8, a_{10}] \oplus \mathbb{Z}_3\{a_4, a_8\} \otimes \mathbb{Z}_3[x_{26}]^+ \oplus \mathbb{Z}_3[a_{10}] \otimes \mathbb{Z}_3[x_{26}]^+; \tag{3.9}$$

$$\begin{aligned}\mathrm{Ker}\ \widetilde{d}'_4 \mid \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{16}b_{18}^2\} &= 0,\\ \mathrm{Coker}\ \widetilde{d}'_4 \mid \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{16}b_{18}^2\} &= \Big(\mathbb{Z}_3[a_8, a_{10}, x_{26}]/x_{26}(a_8, a_{10})\Big)\{y_{26}\}\\ &= (\mathbb{Z}_3[a_8, a_{10}] \oplus \mathbb{Z}_3[x_{26}]^+)\{y_{26}\}.\end{aligned}$$

If we set a submodule

$$\begin{aligned}L = {}& \mathbb{Z}_3[a_8, a_{10}]\{y_{26}\} \oplus \mathbb{Z}_3[x_{26}]^+\{y_{26}\}\\ &\oplus \mathbb{Z}_3[a_4, a_8, a_{10}]\{y_{22}\} \oplus \mathbb{Z}_3[x_{26}]^+\{y_{22}\}\\ &\oplus \mathbb{Z}_3[a_4, a_8, a_{10}]\{y_{20}^2, y_{22}^2, y_{20}y_{22}, y_{58}, y_{60}, y_{76}\}\\ &\oplus \mathbb{Z}_3[a_8, a_{10}]\{y_{20}y_{26}, y_{22}y_{26}, y_{64}\}\\ &\oplus \mathbb{Z}_3[x_{26}]\{a_9, y_{21}, y_{25}, y_{27}, y_{21}a_8, y_{21}a_{10}, y_{25}a_{10}, y_{21}y_{26}\},\end{aligned} \tag{3.10}$$

we obtain that

$$\begin{aligned}\widetilde{E}'_5 = \Big(L &\oplus \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4^2, a_4a_8, a_4a_{10}, a_8^2, a_8a_{10})\\ &\oplus \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4, a_8)\{y_{20}\}\\ &\oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9a_8b_{12}b_{18}^2\} \oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{18}^2\}\Big) \otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}].\end{aligned}$$

To determine $\widetilde{E}'_6$, we need the following

Lemma 3.4. $\widetilde{d}'_5(a_9a_8b_{12}b_{18}^2) = -x_{26}a_{10}^2y_{20}$.

Proof. It follows from (3.6) that

$$d\Big(a_8(a_9b_{12}b_{18}^2 - c_{17}\partial(b_{12}b_{18}^2))\Big) = -x_{26}a_8a_{10}y_{22}.$$

Using the relation $-a_8y_{22} + a_{10}y_{20} = \partial(b_{12}b_{16}b_{18})$ and the formula (3.6), we have that

$$d\Big(a_{10}(a_9b_{12}b_{16}b_{18} - c_{17}\partial(b_{12}b_{16}b_{18}))\Big) = -a_{10}x_{26}(-a_8y_{22} + a_{10}y_{20}).$$

Hence we obtain that

$$d\Big(a_9a_8b_{12}b_{18}^2 - c_{17}a_8\partial(b_{12}b_{18}^2) + a_9a_{10}b_{12}b_{16}b_{18} + c_{17}a_{10}\partial(b_{12}b_{16}b_{18})\Big) = -x_{26}a_{10}^2y_{20},$$

which implies the lemma. □

Therefore the only non trivial differential

$$\widetilde{d_5'} : \mathbb{Z}_3[a_{10}, x_{26}]\{a_9a_8b_{12}b_{18}^2\} \to \Big(\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4, a_8)\Big)\{y_{20}\}$$

is given by $\widetilde{d_5'}(a_9a_8b_{12}b_{18}^2) = -x_{26}a_{10}^2y_{20}$.

Hence we have

$$\begin{aligned} \operatorname{Ker} \widetilde{d_5'} \mid \mathbb{Z}_3[a_{10}, x_{26}]\{a_9a_8b_{12}b_{18}^2\} &= 0, \\ \operatorname{Coker} \widetilde{d_5'} \mid \mathbb{Z}_3[a_{10}, x_{26}]\{a_9a_8b_{12}b_{18}^2\} &= \Big(\mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4, a_8, a_{10}^2)\Big)\{y_{20}\} \\ &= \mathbb{Z}_3[a_4, a_8, a_{10}]\{y_{20}\} \oplus \mathbb{Z}_3[x_{26}]^+\{y_{20}, a_{10}y_{20}\}. \end{aligned}$$

Thus we obtain that

$$\begin{aligned} \widetilde{E_6'} = \Big(L &\oplus \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4^2, a_4a_8, a_4a_{10}, a_8^2, a_8a_{10}) \\ &\oplus \mathbb{Z}_3[a_4, a_8, a_{10}]\{y_{20}\} \oplus \mathbb{Z}_3[x_{26}]^+\{y_{20}, a_{10}y_{20}\} \\ &\qquad\qquad \oplus \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{18}^2\}\Big) \otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}], \end{aligned}$$

where we recall that L is defined at (3.10).

Finally we observe that the following is the only non trivial differential

$$\widetilde{d_6'} : \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{18}^2\} \to \mathbb{Z}_3[a_4, a_8, a_{10}, x_{26}]/x_{26}(a_4^2, a_4a_8, a_4a_{10}, a_8^2, a_8a_{10})$$

given by $\widetilde{d_6'}(a_9b_{18}^2) = -x_{26}a_{10}^2$. Using (3.9), we have the presentation

$$\begin{aligned} \operatorname{Ker} \widetilde{d_6'} \mid \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{18}^2\} &= 0, \\ \operatorname{Coker} \widetilde{d_6'} \mid \mathbb{Z}_3[a_{10}, x_{26}]\{a_9b_{18}^2\} &= \mathbb{Z}_3[a_4, a_8, a_{10}] \oplus \mathbb{Z}_3[x_{26}]^+\{1, a_4, a_8, a_{10}\}. \end{aligned}$$

Thus we obtain that

$$\begin{aligned} \widetilde{E_7'} = \Big(&\mathbb{Z}_3[a_4, a_8, a_{10}]\{1, y_{20}, y_{20}^2, y_{22}, y_{22}^2, y_{20}y_{22}, y_{58}, y_{60}, y_{76}\} \\ &\oplus \mathbb{Z}_3[a_8, a_{10}]\{y_{26}, y_{26}^2, y_{20}y_{26}, y_{22}y_{26}, y_{64}\} \\ &\oplus \mathbb{Z}_3[x_{26}]^+\{1, a_4, a_8, a_{10}, y_{20}, a_{10}y_{20}, y_{22}, y_{26}\} \\ &\oplus \mathbb{Z}_3[x_{26}]\{a_9, y_{21}, y_{25}, y_{27}, y_{21}a_8, y_{21}a_{10}, y_{25}a_{10}, y_{21}y_{26}\}\Big) \\ &\otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}]. \end{aligned}$$

Since all the generators a_i, x_j, y_k chosen in Lemma 3.2 can be seen to be cocycles by direct calculation in the differential algebra $(\overline{V}, d)$, we see that $\widetilde{E_7'} = \widetilde{E_\infty'} = E_7$ and $E_7 = E_\infty$. Thus we have proved Theorem 2.1.

4. Proof of Theorem 2.3

Using the presentation of Theorem 2.1, we can list up all the relations, which are calculated in the complex $(\overline{V}, d)$ defined in Section 3. The calculation is usually straightforward, although it becomes more systematic when using the auxiliary derivation ∂ defined in (3.3).

First we need two lemmas.

Lemma 4.1. *We have that*

$$a_9\partial^2 Q = y_{21}\partial^2 Q = y_{25}\partial^2 Q = y_{27}\partial^2 Q = x_{26}\partial^2 Q = 0,$$

where Q is an element of $\mathbb{Z}_3[a_4, a_8, a_{10}, b_{12}, b_{16}, b_{18}]$.

Proof. $a_9\partial^2 Q = d(\partial Q)$, $\quad y_{21}\partial^2 Q = d(a_4 Q + b_{12}\partial Q)$,
$y_{25}\partial^2 Q = d(a_8 Q + b_{16}\partial Q)$, $\quad y_{27}\partial^2 Q = d(a_{10} Q + b_{18}\partial Q)$.
The last equation was proved in Lemma 3.3. ☐

Lemma 4.2. *The ∂^2-image is generated over $\mathbb{Z}_3[a_4, a_8, a_{10}, x_{36}, x_{48}, x_{54}]$ by the elements in the table* (4.0) *in the next page.*

Proof of Theorem 2.3.B One can show the relations in i) by expressing the elements in terms of a_i and b_j and those in ii) by case by case checking as follows:

$$a_9^2 = d(c_{17}), \quad y_{21}^2 = d(c_{17}b_{12}^2), \quad y_{25}^2 = d(c_{17}b_{16}^2), \quad y_{27}^2 = d(c_{17}b_{18}^2),$$
$$a_9y_{21} + x_{26}a_4 = d(c_{17}b_{12}), \quad a_9y_{25} + x_{26}a_8 = d(c_{17}b_{16}), \quad a_9y_{27} + x_{26}a_{10} = d(c_{17}b_{18}),$$
$$y_{21}y_{25} + x_{26}y_{20} = d(c_{17}b_{12}b_{16}), \quad y_{21}y_{27} - x_{26}y_{22} = d(c_{17}b_{12}b_{18}), \quad y_{25}y_{27} - x_{26}y_{26} = d(c_{17}b_{16}b_{18}).$$

The first five relations in iii) are immediate from Lemmas 4.1 and 3.3. One can show the remaining relations by appealing to the relation $d(Q) = a_9\partial Q + c_{17}\partial^2 Q$ for $Q \in \mathbb{Z}_3[a_i, b_j]$ and the table in Lemma 4.2:

$$a_9a_4 = d(b_{12}), \quad a_9a_8 = d(b_{16}), \quad a_9a_{10} = d(b_{18}),$$
$$y_{21}a_4 = d(b_{12}^2), \quad y_{25}a_8 = d(b_{16}^2), \quad y_{27}a_{10} = d(b_{18}^2),$$
$$y_{21}a_8 + a_9y_{20} = y_{25}a_4 - a_9y_{20} = d(b_{12}b_{16}),$$
$$y_{21}a_{10} - a_9y_{22} = y_{27}a_4 + a_9y_{22} = d(b_{12}b_{18}),$$
$$y_{25}a_{10} - a_9y_{26} = y_{27}a_8 + a_9y_{26} = d(b_{16}b_{18}),$$
$$y_{21}y_{20} = d(-b_{12}^2b_{16}), \quad y_{25}y_{20} = d(-b_{12}b_{16}^2),$$
$$y_{21}y_{22} = d(-b_{12}^2b_{18}), \quad y_{27}y_{22} = d(-b_{12}b_{18}^2),$$
$$y_{25}y_{26} = d(-b_{16}^2b_{18}), \quad y_{27}y_{26} = d(-b_{16}b_{18}^2),$$
$$y_{27}y_{20} - y_{25}y_{22} = y_{25}y_{22} + y_{21}y_{26} = -y_{27}y_{20} - y_{21}y_{26} = d(-b_{12}b_{16}b_{18}).$$

Proof of Theorem 2.3.A First we have to show that $D \otimes \mathbb{Z}_3[x_{36}, x_{48}, x_{54}]$ is an ideal of $\mathrm{Cotor}_{H^*(E_6)}(\mathbb{Z}_3, \mathbb{Z}_3)$. We denote by S a subalgebra

$$\mathbb{Z}_3[a_4, a_8, a_{10}, b_{12}, b_{16}, b_{18}]$$

of $\overline{V}$ defined in Section 3. By Lemma 3.1 we obtain a module isomorphism

$$\overline{V} \cong T(a_9, c_{17}) \otimes S = T(a_9, c_{17})^+ \otimes S \oplus S.$$

Table (4.0).

Q	∂Q	$\partial^2 Q$
b_{12}	$-a_4$	0
b_{16}	$-a_8$	0
b_{18}	$-a_{10}$	0
b_{12}^2	a_4b_{12}	$-a_4^2$
b_{16}^2	a_8b_{16}	$-a_8^2$
b_{18}^2	$a_{10}b_{18}$	$-a_{10}^2$
$b_{12}b_{16}$	$-a_4b_{16}-a_8b_{12}$	$-a_4a_8$
$b_{12}b_{18}$	$-a_4b_{18}-a_{10}b_{12}$	$-a_4a_{10}$
$b_{16}b_{18}$	$-a_8b_{18}-a_{10}b_{16}$	$-a_8a_{10}$
$b_{12}^2b_{16}$	$a_4b_{12}b_{16}-a_8b_{12}^2$	$-a_4y_{20} = -a_4^2b_{16}+a_4a_8b_{12}$
$b_{12}b_{16}^2$	$-a_4b_{16}^2+a_8b_{12}b_{16}$	$a_8y_{20} = a_4a_8b_{16}-a_8^2b_{12}$
$b_{12}^2b_{18}$	$a_4b_{12}b_{18}-a_{10}b_{12}^2$	$-a_4y_{22} = -a_4^2b_{18}+a_4a_{10}b_{12}$
$b_{12}b_{18}^2$	$-a_4b_{18}^2+a_{10}b_{12}b_{18}$	$a_{10}y_{22} = a_4a_{10}b_{18}-a_{10}^2b_{12}$
$b_{16}^2b_{18}$	$a_8b_{16}b_{18}-a_{10}b_{16}^2$	$-a_8y_{26} = -a_8^2b_{18}+a_8a_{10}b_{16}$
$b_{16}b_{18}^2$	$-a_8b_{18}^2+a_{10}b_{16}b_{18}$	$a_{10}y_{26} = a_8a_{10}b_{18}-a_{10}^2b_{16}$
$b_{12}b_{16}b_{18}$	$-a_4b_{16}b_{18}-a_8b_{12}b_{18}-a_{10}b_{12}b_{16}$	$-a_4y_{26}+a_{10}y_{20} = -a_8y_{22}-a_{10}y_{20} = a_4y_{26}+a_8y_{22}$
		$= -a_4a_8b_{18}-a_4a_{10}b_{16}-a_8a_{10}b_{12}$
$b_{12}^2b_{16}^2$	$a_4b_{12}b_{16}^2+a_8b_{12}^2b_{16}$	$-y_{20}^2 = -a_4^2b_{16}^2-a_4a_8b_{12}b_{16}-a_8^2b_{12}^2$
$b_{12}^2b_{18}^2$	$a_4b_{12}b_{18}^2+a_{10}b_{12}^2b_{18}$	$-y_{22}^2 = -a_4^2b_{18}^2-a_4a_{10}b_{12}b_{18}-a_{10}^2b_{12}^2$
$b_{16}^2b_{18}^2$	$a_8b_{16}b_{18}^2+a_{10}b_{16}^2b_{18}$	$-y_{26}^2 = -a_8^2b_{18}^2-a_8a_{10}b_{16}b_{18}-a_{10}^2b_{16}^2$
$b_{12}^2b_{16}b_{18}$	$a_4b_{12}b_{16}b_{18}-a_8b_{12}^2b_{18}-a_{10}b_{12}^2b_{16}$	$-y_{20}y_{22} = -a_4^2b_{16}b_{18}+a_4a_8b_{12}b_{18}+a_4a_{10}b_{12}b_{16}-a_8a_{10}b_{12}^2$
$b_{12}b_{16}^2b_{18}$	$-a_4b_{16}^2b_{18}+a_8b_{12}b_{16}b_{18}-a_{10}b_{12}b_{16}^2$	$y_{20}y_{26} = a_4a_8b_{16}b_{18}-a_4a_{10}b_{16}^2-a_8^2b_{12}b_{18}+a_8a_{10}b_{12}b_{16}$
$b_{12}b_{16}b_{18}^2$	$-a_4b_{16}b_{18}^2-a_8b_{12}b_{18}^2+a_{10}b_{12}b_{16}b_{18}$	$-y_{22}y_{26} = -a_4a_8b_{18}^2+a_4a_{10}b_{16}b_{18}+a_8a_{10}b_{12}b_{18}-a_{10}^2b_{12}b_{16}$
$b_{12}^2b_{16}^2b_{18}$	$a_4b_{12}b_{16}^2b_{18}+a_8b_{12}^2b_{16}b_{18}-a_{10}b_{12}^2b_{16}^2$	$y_{58} = -a_4^2b_{16}^2b_{18}-a_4a_8b_{12}b_{16}b_{18}+a_4a_{10}b_{12}b_{16}^2$
		$-a_8^2b_{12}^2b_{18}+a_8a_{10}b_{12}^2b_{16}$
$b_{12}^2b_{16}b_{18}^2$	$a_4b_{12}b_{16}b_{18}^2-a_8b_{12}^2b_{18}^2+a_{10}b_{12}^2b_{16}b_{18}$	$y_{60} = -a_4^2b_{16}b_{18}^2+a_4a_8b_{12}b_{18}^2-a_4a_{10}b_{12}b_{16}b_{18}$
		$+a_8a_{10}b_{12}^2b_{18}-a_{10}^2b_{12}^2b_{16}$
$b_{12}b_{16}^2b_{18}^2$	$-a_4b_{16}^2b_{18}^2+a_8b_{12}b_{16}b_{18}^2+a_{10}b_{12}b_{16}^2b_{18}$	$y_{64} = a_4a_8b_{16}b_{18}^2+a_4a_{10}b_{16}^2b_{18}-a_8^2b_{12}b_{18}^2$
		$-a_8a_{10}b_{12}b_{16}b_{18}-a_{10}^2b_{12}b_{16}^2$
$b_{12}^2b_{16}^2b_{18}^2$	$a_4b_{12}b_{16}^2b_{18}^2+a_8b_{12}^2b_{16}b_{18}^2+a_{10}b_{12}^2b_{16}^2b_{18}$	$y_{76} = -a_4^2b_{16}^2b_{18}^2-a_4a_8b_{12}b_{16}b_{18}^2-a_4a_{10}b_{12}b_{16}^2b_{18}$
		$-a_8^2b_{12}^2b_{18}^2-a_8a_{10}b_{12}^2b_{16}b_{18}-a_{10}^2b_{12}^2b_{16}^2$

Now we will show that $T(a_9,c_{17})^+\otimes S$ is a two sided ideal of $\overline{V}$. Recall the relations in $\overline{V}$:

$$\begin{aligned} b_ja_9 &= a_9b_j + c_{17}a_{j-8} && \text{for}\quad j=12,16,18;\\ a_ja_9 &= a_9a_j && \text{for}\quad j=4,8,10;\\ Qc_{17} &= c_{17}Q && \text{for}\quad Q\in S, \end{aligned}$$

and also observe that

$$QT(a_9,c_{17})^+\otimes S\subset T(a_9,c_{17})^+\quad\text{for}\quad Q\in S$$

which is obtained by the repeated use of the above relations. Thus $T(a_9,c_{17})^+\otimes S$ is a two sided ideal of $\overline{V}$. Let $Z(\overline{V})$ and $B(\overline{V})$ be the sub-groups consisting of cocycles and coboundaries in $\overline{V}$ respectively. Thus we have

$$\begin{aligned} Z(\overline{V}) &= \Big((T(a_9,c_{17})^+\otimes S)\cap Z(\overline{V})\Big)\oplus S\cap Z(\overline{V}),\\ B(\overline{V}) &= \Big((T(a_9,c_{17})^+\otimes S)\cap B(\overline{V})\Big)\oplus S\cap B(\overline{V}). \end{aligned}$$

Using the fact that $T(a_9,c_{17})^+\otimes S$ is a two sided ideal of $\overline{V}$, we can show that

$$(T(a_9,c_{17})^+\otimes S)\cap Z(\overline{V})\text{ and }(T(a_9,c_{17})^+\otimes S)\cap Z(\overline{V})$$

are both two sided ideals of $Z(\overline{V})$. Hence

$$\frac{(T(a_9,c_{17})^+\otimes S)\cap Z(\overline{V})}{(T(a_9,c_{17})^+\otimes S)\cap B(\overline{V})}$$

is an ideal of $H^*(\overline{V})=\mathrm{Cotor}_{H^*(E_6)}(\mathbb{Z}_3,\mathbb{Z}_3)$.

Here we have

$$D\otimes\mathbb{Z}_3[x_{36},x_{48},x_{54}]=\frac{(T(a_9,c_{17})^+\otimes S)\cap Z(\overline{V})}{(T(a_9,c_{17})^+\otimes S)\cap B(\overline{V})}\tag{4.1}$$

$$C\otimes\mathbb{Z}_3[x_{36},x_{48},x_{54}]=\frac{S\cap Z(\overline{V})}{S\cap B(\overline{V})}\tag{4.2}$$

In particular, it follows from (4.1) that $D\otimes\mathbb{Z}_3[x_{36},x_{48},x_{54}]$ is an ideal of $\mathrm{Cotor}_{H^*(E_6)}(\mathbb{Z}_3,\mathbb{Z}_3)$.

The remaining part is to check the splitness. By the definition, it means that a relation in $C\otimes\mathbb{Z}_3[x_{36},x_{48},x_{54}]$ is also that of $\mathrm{Cotor}_{H^*(E_6)}(\mathbb{Z}_3,\mathbb{Z}_3)$. This fact is a direct consequence of checking the relations stated in Theorem 2.3.B.

5. The May spectral sequence

In this section, we review our calculations from the view point of the May spectral sequence [May1, 2]. For the sake of convenience, we recall the construction so that it fits to our situation. Let us recall from [KM] the following

Theorem 5.1. *As an algebra, we have*

$$H^*(E_6) \cong \mathbb{Z}_3[x_8]/(x_8^3) \otimes \Lambda(x_3, x_7, x_9, x_{11}, x_{15}, x_{17}),$$

where the reduced diagonal map induced from the multiplication $\widetilde{\varphi} : \widetilde{H}^*(E_6) \to \widetilde{H}^*(E_6) \otimes \widetilde{H}^*(E_6)$ *is given by*

$$\begin{aligned} &\widetilde{\varphi}(x_j) = x_8 \otimes x_{j-8} && \textit{for} \quad j = 11, 15, 17, \\ &\widetilde{\varphi}(x_j) = 0 && \textit{otherwise.} \end{aligned}$$

Let $A = H^*(E_6)$ and I the augmented ideal generated by the elements x_3, x_7, x_9, x_{11}, x_{15}, x_{17}. Let $C^*_A(\mathbb{Z}_3, \mathbb{Z}_3)$ be the cobar complex [May1, 2] and $[x_1|\cdots|x_n]$ denote an element of $C^i_A(\mathbb{Z}_3, \mathbb{Z}_3) = I^{\otimes i}$ and $C^0_A(\mathbb{Z}_3, \mathbb{Z}_3) = [\ ] \cong \mathbb{Z}_3$. Then we can define a weight v of $[x_1|\cdots|x_n]$ by

$$\min\{n_1 + \cdots + n_i \mid x_j \in I^{n_j},\ j = 1, \cdots, i\}\,;$$

let $F^pC^*_A(\mathbb{Z}_3, \mathbb{Z}_3)$ be the ideal generated by the elements $[x_1|\cdots|x_n]$ where $1 \le i$ such that $v([x_1|\cdots|x_n]) \ge p$.

The construction of an injective resolution (see [SI] and [MS]) induces a surjection p from $C^*_A(\mathbb{Z}_3, \mathbb{Z}_3)$ to $\overline{V}$ defined in Section 3; more concretely it is described as $p([x_3]) = a_4$, $p([x_7]) = a_8$, $p([x_8]) = a_9$, $p([x_8^2]) = c_{17}$, $p([x_{11}]) = b_{12}$, $p([x_{15}]) = b_{16}$, $p([x_{17}]) = b_{18}$. We see that $w([x_8^2]) = 2$ and $w([x_i]) = 1$ for the other elements. Since $w(a_9b_j) = w(b_ja_9 - c_{17}a_{j-8}) = 2$ for $j = 12, 16, 18$, we can introduce the filtration $F^p\overline{V}$ in $\overline{V}$ by $p(F^pC^*_A(\mathbb{Z}_3, \mathbb{Z}_3))$. We also use the same letter w for the weight of an element of $\overline{V}$. Then we have $w(c_{17}) = 2$ and $w(a_i) = w(b_i) = 1$. We denote by $\{E_p(\overline{V}), d_p\}$ the spectral sequence of $\overline{V}$ induced from the above filtration.

Lemma 5.2. (i) *We have*

$$E_1(\overline{V}) = \mathbb{Z}_3[x_{26}] \otimes \mathbb{Z}_3[a_4, a_8, a_{10}] \otimes \Lambda(a_9) \otimes \mathbb{Z}_3[b_{12}, b_{16}, b_{18}],$$

where $w(a_4) = w(a_8) = w(a_{10}) = w(a_9) = w(b_{12}) = w(b_{16}) = w(b_{18}) = 1$ *and* $w(x_{26}) = 3$.

(ii) *After the* $E_p(\overline{V})$*-term for* $p \ge 1$*, the spectral sequence coincides with the May spectral sequence which converges to* $\mathrm{Cotor}_{H^*(E_6)}(\mathbb{Z}_3, \mathbb{Z}_3)$.

Proof. Since the only non trivial differential is given by $d_0(c_{17}) = a_9^2$, the argument to obtain $E_1(\overline{V})$ is the same as Section 3. The rest of the lemma follows from the definitions of the May spectral sequence and the filtration. □

The next lemma follows from the proof of Lemma 3.2.

Lemma 5.3. $E_2(\overline{V})$ *is obtained from the differential*

$$d_1(b_{12}) = -a_4a_9, \quad d_1(b_{16}) = -a_8a_9, \quad d_1(b_{18}) = -a_{10}a_9.$$

Hence $E_2(\overline{V})$ *is given in the formula of Lemma* 3.2

At this point, the meaning of Lemma 3.7 given by [MS] becomes very clear.

Proposition 5.4. *$E_3(\overline{V})$ is obtained from the differential*

$$d_2(a_9Q) = -x_{26}\partial^2(Q) \quad \textit{for} \quad Q \in \mathbb{Z}_3[a_4, a_8, a_{10}, b_{12}, b_{16}, b_{18}].$$

Furthermore, the May spectral sequence collapses at $E_3(\overline{V})$.

Proof. By Lemma 3.3, we have

$$x_{26}\partial^2(-Q) = d(a_9Q + c_{17}\partial Q).$$

For the elements indicated from (3) to (7) at Lemma 3.2, it is easily shown that

$$\begin{aligned} w(a_9Q + c_{17}\partial Q) &= \min\{w(a_9Q), w(c_{17}\partial Q)\} = 1 + w(Q), \\ w(x_{26}\partial^2(-Q)) &= w(x_{26}) + w(\partial^2(-Q)) = 3 + w(Q). \end{aligned}$$

So we obtain the assertion since $a_9Q + c_{17}\partial Q$ is represented by a_9Q at $E_2(V)$. The other statement follows from the calculation of E_7 in Section 3. ☐

Remark 5.5 The argument in this section is hinted by the second work of Evens and Siegel [ES]. They have given an example of an ungraded Hopf algebra with the type P such that the May spectral sequence does not collapse at the E_2-term. Corresponding to their example is ours in a graded sense. The calculation of [MS] for other exceptional Lie groups for odd prime seems probably to imply that the May spectral sequence converging to $\mathrm{Cotor}_{H^*(G)}(\mathbb{Z}_p, \mathbb{Z}_p)$ collapses at the E_3-level but does not at the E_2-level possibly except the case $(E_8, 3)$.

REFERENCES

[ES] L. Evens and S. F. Siegel : Cohomology of p-restricted Lie algebra, preprint.

[IS] A. Iwai and N. Shimada : A remark on resolutions for Hopf algebras, Publ. RIMS of Kyoto Univ., **1**(1966), 187–198.

[KM] A. Kono and M. Mimura : Cohomology mod 3 of the classifying space of the Lie group E_6, Math. Scand, **46**(1980) 223–235.

[KMS] A. Kono, M. Mimura and N. Shimada : Cohomology of classifying space of certain associative H-space, J. Math. Kyoto Univ., **15**(1975) 607–617.

[May1] J. P. May : The cohomology of restricted Lie algebras and of Hopf algebras; application to the Steenrod algebra, Princeton Univ., mimeographed note, 1964.

[May2] J. P. May : The cohomology of restricted Lie algebras and of Hopf algebras, Bull. AMS, **71**(1965), 372–377.

[M] M. Mimura : Hopf Spaces, Kinokuniya, **26** (1986) (in Japanese).

[MS] M. Mimura and Y. Sambe : On the cohomology mod p of the classifying spaces of the exceptional Lie groups, I, J. Math. Kyoto Univ.,**19**(1979), 553–581.

[MST1] M. Mimura, Y. Sambe and M. Tezuka : Cohomology mod 3 of the classifying spaces of the exceptional Lie group of type E_6, I, (1986/1991), preprint.

[MST2] M. Mimura, Y. Sambe and M. Tezuka : Some remarks on the mod 3 cohomology of the classifying space of the exceptional Lie group E_6, Proceedings of Workshop in Pure Mathematics, Part III, 17(1998), 139 - 159.

[RS] M. Rothenberg and N. E. Steenrod : The cohomology of classifying spaces of H-spaces, Bull. AMS, **71**(1965), 872–875.

[SI] N. Shimada and A. Iwai : On the cohomology of some Hopf algebra, Nagoya Math. J., **30**(1967), 103–111.

[T] H. Toda : Cohomology of classifying space of exceptional Lie groups, Manifolds Tokyo 1973, 265–271.

DEPARTMENT OF MATHEMATICS FACULTY OF SCIENCE OKAYAMA UNIVERSITY; 3-1-1 TSUSHIMA-NAKA, OKAYAMA 700-8530, JAPAN
E-mail address: `mimura@math.okayama-u.ac.jp`

SCHOOL OF BUSINESS ADMINISTRATION SENSHU UNIVERSITY; 2-1-1 TAMA-KU, KAWASAKI 214-8580, JAPAN
E-mail address: `thm0166@isc.senshu-u.ac.jp`

DEPARTMENT OF MATHEMATICAL SCIENCES FACULTY OF SCIENCE UNIVERSITY OF RYUKYUS; NISHIHARA-CHO, OKINAWA 903-0213, JAPAN
E-mail address: `tez@math.u-ryukyu.ac.jp`